

Name: \_\_\_\_\_

Pid: \_\_\_\_\_

1. (10 points) Give a natural deduction derivation of  $\exists x (A(x) \vee B(x))$  from  $\exists x A(x) \vee \exists x B(x)$ .

2. (10 points) Let us consider the following formulas on the variables from the set  $\{x_0, \dots, x_n\}$ .
1. The formula  $I_n$  is equal to  $x_0$ .
  2. The formula  $S_{n,i}$  is equal to  $x_{i-1} \implies x_i$ .
  3. The formula  $T_n$  is equal to  $x_n$ .

Show that there is a natural deduction derivation of  $T_n$  from  $I_n \wedge \bigwedge_{i=1}^n S_{n,i}$ .

3. (10 points) Let  $\phi = \bigvee_{i=1}^m \lambda_i$  be a clause; we say that the width of the clause is equal to  $m$ . Let  $\phi = \bigwedge_{i=1}^{\ell} \chi_i$  be a formula in CNF; we say that the width of  $\phi$  is equal to the maximal width of  $\chi_i$  for  $i \in [\ell]$ .

Let  $m_n : \{T, F\}^n \rightarrow \{T, F\}$  such that  $m_n(x_1, \dots, x_n) = T$  iff the number of elements in the set  $\{i : x_i = T\}$  is divisible by 3.

Show that any CNF representation of  $m_n$  has width at least  $n - 2$ .

4. (10 points) Let  $A\Delta B = (A \cup B) \setminus (A \cap B)$ ; we say that  $A\Delta B$  is the symmetric difference of  $A$  and  $B$ . Let  $\Omega$ , and  $A_1, \dots, A_n \subseteq \Omega$  be some sets. We say that  $\Delta_{i=1}^1 A_i = A_1$  and  $\Delta_{i=1}^{k+1} A_i = (\Delta_{i=1}^k A_i) \Delta A_{k+1}$ . Show that

$$\Delta_{i=1}^n A_i = \{x \in \Omega : x \in A_i \text{ for odd number of } i \in [n]\}.$$

5. (10 points) Let  $\mathcal{S}$  be a signature with two predicate symbols  $=$  and  $S$  such that the first is binary and the last is ternary.

Let us consider the structure  $\mathfrak{M}$  such that it corresponds to the points on a two-dimensional plane,  $=$  is a standard equality, and  $S(x, y, z)$  is true iff  $|xz| = |yz|$ .

Let  $R$  be a relation such that  $(A, B, C) \in R$  iff  $A$ ,  $B$ , and  $C$  lay on the same line. Show that  $R$  is representable in  $\mathfrak{M}$ .

6. (10 points) Let us define the set  $S$  defined as follows:

- $3 \in S$  and
- if  $x \in S$  and  $y \in S$ , then  $(x + y) \in S$ .

Show that  $S = \{3k : k \in \mathbb{N}\}$ .

7. (10 points) Let  $f, g_1, \dots, g_n : \mathbb{R}^\ell \rightarrow \mathbb{R}$ . We say that the equation  $f(x) = 0$  can be derived from the equations  $g_1(x) = 0, \dots, g_n(x) = 0$  iff there is a sequence of functions  $h_1, \dots, h_m : \mathbb{R}^\ell \rightarrow \mathbb{R}$  such that  $h_m = f$  and for each  $i \in [m]$ ,

- either  $h_i$  is equal to  $g_j$  for some  $j \in [n]$ , or
- $h_i = h_j + h_k$  for some  $1 \leq j, k < i$ , or
- $h_i = c \cdot h_j$  for some  $1 \leq j < i$  and some  $c \in \mathbb{R}$ .

Show that if the equation  $f(x) = 0$  can be derived from the equations  $g_1(x) = 0, \dots, g_n(x) = 0$ , then for any  $v \in \mathbb{R}^\ell$ ,  $f(v) = 0$  provided that  $g_1(v) = \dots = g_n(v) = 0$ .